

Hidden Chemistry of Non-Enzymatic Acyl Transfer in Cellular Systems

Mahmood Khalid Hassoon ^{1,*} and Maan Jowad Zraow ²

¹ Department of Chemistry, College of Education for Pure Sciences, University of Kirkuk, Kirkuk, Iraq.

² Department of Forensic Evidences, College of Science, University of Kirkuk, Kirkuk, Iraq.

Open Access Research Journal of Life Sciences, 2026, 11(02), 096–105

Publication history: Received on 27 February 2026; revised on 03 April 2026; accepted on 06 April 2026

Article DOI: <https://doi.org/10.53022/oarjls.2026.11.2.0007>

Abstract

Non-enzymatic protein modifications are a type of post-translational modification that occurs as a result of a direct chemical reaction between active acyl donors, such as acyl-CoA, and nucleophilic groups in proteins without enzymatic intervention. This phenomenon has received increasing attention due to its potential role in linking the metabolic state of the cell to the functions of proteins. Despite its importance, the mechanistic understanding of non-enzymatic fluid interactions is still limited, especially with regard to how environmental factors within the cell affect their behavior. Most previous studies have focused on descriptive aspects or enzymatic mechanisms, while the kinetics and specificity of these reactions under physiological conditions are still not sufficiently clear. Therefore, this research aims to develop an integrative conceptual framework that links intracellular conditions, such as pH and acyl-CoA concentration, and between the dynamics and selectivity of the interaction. The research analyzes how changes in the cellular environment affect nucleophilicity, reaction rates, and the likelihood of modification at specific sites on proteins. The results suggest that non-enzymatic questions are not entirely random, but are influenced by the interaction of chemical agents and local protein environments that lead to preferential patterns of modulation under certain metabolic conditions. The results also showed that increased acyl-CoA availability and pH change are key factors in enhancing reactivity and influencing the distribution of modified sites. Overall, this research demonstrates that non-enzymatic questions are a process dependent on cellular context and reflect the interaction between metabolic status and protein chemistry, providing a more comprehensive understanding of their functional role within biosystems.

Keywords: Hidden chemistry, Non-enzymatic acyl transport, Non-enzymatic fluids, Cellular systems, Intracellular chemical reactions

1 Introduction

Post-translational modifications (PTMs) are essential biological mechanisms that contribute to the expansion of the functional diversity of proteins after their translation, as they play a pivotal role in regulating enzymatic activity, protein stability, molecular interactions, and cellular signaling pathways (Walsh et al., 2005). Among these modifications, protein acylation represents an important pattern that involves the addition of acyl groups to specific amino acid residues, leading to substantial changes in the physical, chemical, and functional properties of proteins (Sabari et al., 2017).

Protein question interactions can be classified into two main types: enzymatic acylation and non-enzymatic acylation. Enzymatic questions rely on specialized enzymes such as acyltransferases that transport the acyl group from active donors such as acyl-CoA to specific sites on proteins with a high degree of selectivity and regulation (Marmorstein & Zhou, 2014). In contrast, non-enzymatic questions occur as a result of a direct chemical reaction between acyl-CoA molecules and nucleophilic groups in proteins, without enzymatic intervention, making them more related to the chemical properties of the cellular medium such as pH and metabolite concentration (Hirschey et al., 2015).

* Corresponding author: Mahmood Khalid Hassoon

These interactions are particularly important due to their close association with metabolic status within the cell, as acyl-CoA levels reflect changes in metabolic pathways, making protein questions a meeting point between metabolic regulation and protein modification (Lin et al., 2012). 2014).

Despite significant advances in the study of protein questions, most previous research has focused primarily on descriptive aspects or enzymatic mechanisms, while the mechanistic and quantitative understanding of non-enzymatic interactions remains limited (Weinert et al., 2013). The effect of intracellular factors—such as pH, acyl-CoA concentrations, and protein structure—on both kinetics and specificity has not been studied in an integrative and comprehensive manner (Huang et al., 2018).

The research gap is the absence of an analytical framework that links the chemical properties of acyl-CoA molecules, the kinetics and selectivity of reactions, within the context of the cellular environment. In this sense, this study aims to develop a conceptual and analytical model that explains the interactions of non-enzymatic questions by integrating three main axes: kinetics, selectivity, and intracellular conditions.

This raises the following research question: How do pH and acyl-CoA concentration within an integrative framework affect the kinetics and selectivity of intracellular non-enzymatic fluid reactions?

1.1 Research Objective

The aim of this research is to analyze and interpret the effect of pH and acyl-CoA concentration within a conceptual integrative framework, to understand their role in determining kinetics and specificity of intracellular non-enzymatic fluid interactions.

1.2 Importance of Research

Scientific importance lies in providing an analytical framework that explains non-enzymatic reactions in an integrative way that combines biochemistry and kinetics, rather than the traditional descriptive proposition. The study also contributes to clarifying the relationship between protein modifications and the metabolic state of the cell. In practice, the results help guide experimental studies and improve understanding of protein changes associated with metabolic diseases, supporting the development of more accurate diagnostic and research applications.

1.3 Research Opportunity

Changes in pH (pH) and acyl-CoA concentration, within an integrative reactive framework, coherently regulate the kinetics and selectivity of intracellular non-enzymatic fluid reactions, so that both the ionization state and the availability of the substrate determine the level and efficiency of the reaction.

1.4 Research methodology

This study relies on a narrative review approach to analyze the literature related to the transmission of non-enzymatic acyl groups and their relationship to intracellular conditions, where studies were collected from scientific databases such as PubMed , Scopus, and Web of Science using relevant keywords, and then analyzed qualitatively to derive patterns and conceptual relationships between different variables and build an integrated analytical framework.

2 Theoretical Framework

2.1 Chemical basis for the transmission of non-enzymatic acyl groups

Acyl-CoA molecules are highly reactive biomolecules due to their high degradation potential compared to traditional ester bonds, which gives them a high ability to transport the acyl group and participate in bioreactions within the cell. This property is key to its central role within intermediate metabolism, as it serves as a crossover point between energy production pathways and the regulation of post-translational modifications of proteins (Nelson & Cox, 2021; Berg, Tymoczko, & Stryer, 2019).

Non-enzymatic liquefied reactions arise as a result of the direct interaction between the acyl-CoA acyl group and the nucleophilic groups found on amino acid residues in proteins, especially the amine groups in lysine and the thiol groups in cysteine. This process takes place through the nucleophilic attack mechanism, in which the nucleophilic atom attacks the carbonyl carbon in the thioester bond, resulting in the formation of a new acyl bond with the release of the CoA molecule as an outgoing product (Voet & Voet, 2016; Berg et al., 2019).

The susceptibility of this reaction is influenced by several chemical properties, most notably the high electrophilicity of carbonyl carbon in the thioester bond, which increases its susceptibility to nucleophilic attack. Also, the low stability of the thioester bond compared to other ester bonds makes it more susceptible to degradation and reaction, which explains its high chemical activity (Nelson & Cox, 2021). The spatial structure of the protein plays a crucial role in determining the accessibility of reactive sites, as the tertiary structure can limit the exposure of amino acid residues to nucleophiles, influencing the rate and location of the reaction (Alberts et al., 2015). Physiological conditions within the cell, such as pH and metabolite concentration, affect the ionization state of functional groups, and thus the efficiency and speed of the reaction (Lehninger Principles of Biochemistry, Nelson & Cox, 2021).

Non-enzymatic liquefied reactions do not depend only on the presence of reactants, but their efficiency is determined by a complex interaction between the chemical properties of acyl-CoA, the amino acid properties of proteins, as well as the influence of the cellular environment that acts as a regulatory factor that regulates these reactions and determines their functional outcomes (Voet & Voet, 2016).

2.2 Kinetics of non-enzymatic fluid reactions

The chemical kinetics of non-enzymatic reactions are key to understanding how these reactions occur within the cellular environment, as reaction rates determine the extent to which protein modifications accumulate and their functional effect. In this context, these reactions are described as relying on direct chemical reactions between molecules without enzymatic intervention, making the reaction rate related to the physical and chemical properties of the reactants and their surrounding environment (Nelson & Cox, 2021).

The rates of non-enzymatic reactions are based on conventional laws of chemical kinetics, as they can be expressed through the rate constant (k) which determines the speed of formation of the product relative to the concentrations of reactants. Unlike enzymatic reactions, which often follow the Michaelis–Menten kinetics due to the presence of a catalytic active site, non-enzymatic reactions do not possess this type of regulation and are therefore more directly sensitive to the concentrations of reactants and the conditions of the medium (Berg, Tymoczko, & Stryer, 2019).

When comparing non-enzymatic reactions with their enzymatic counterparts, it is clear that enzymatic reactions are characterized by a significant increase in the rate of reaction due to reduced activation energy through enzymatic catalysis, as well as a high degree of selectivity towards substrates. In contrast, non-enzymatic reactions are relatively slower, but may become physiologically important when high concentrations of acyl-CoA are available or under certain cellular conditions that promote the reaction (Voet & Voet, 2016). The absence of an enzymatic active site also means that selectivity in non-enzymatic reactions depends primarily on the structural accessibility of amino residues rather than specific molecular recognition.

The concentration of reactants is one of the most important factors influencing the reaction rate, as a high concentration of acyl-CoA increases the probability of collision between the reacting molecules, thus increasing the rate of non-enzymatic reactions according to the principles of collision theory (Atkins & de Paula, 2014). The concentration of target proteins, especially those containing exposed lysine or cysteine residues, also plays a role in determining the locations and likelihood of the reaction occurring.

Time affects the accumulation of reaction products, as non-enzymatic reactions may be relatively slow, but over time they can lead to noticeable levels of protein modification, especially in environments where the rate of removal or repair of these modifications is limited (Alberts et al., 2015). Therefore, the interaction is understood not only in terms of its instantaneous velocity, but also in terms of the dynamics of accumulation over time within the cell.

Temperature is also a factor in the kinetic dynamics of the reaction, as increasing the temperature increases the kinetic energy of the molecules, which enhances the rate of effective collisions and increases the speed of the reaction according to the Arrhenius equation. However, the physiological effect of intracellular temperature is limited within a narrow range, but may become significant in experimental or abnormal conditions (Voet & Voet, 2016).

It is clear that the kinetics of non-enzymatic reactions are determined by a complex interaction between several variables, including reactant concentrations, time, temperature, and the properties of the chemical medium. Although these reactions are simple compared to enzymatic reactions in terms of mechanism, their kinetic behavior within the cell reflects a dynamic system that depends on the balance between these factors, making their understanding requires an integrative analysis that combines physical chemistry and cellular biology (Nelson & Cox, 2021; Alberts et al., 2015).

2.3 Selectivity: Is it really random?

The question of specificity in non-enzymatic reaction reactions is central to understanding the extent to which this phenomenon is organized within cells, as it raises a fundamental question of whether these modifications occur entirely randomly or are subject to chemical and structural constraints that make them somewhat selective. Recent evidence suggests that these reactions, in the absence of enzymatic catalysis, do not occur entirely randomly, but are influenced by several factors that determine the sites most susceptible to the occurrence of the reactions (Walsh & Mohr, 2011).

In terms of amino acid selectivity, studies show that lysine residues are one of the most common sites of non-enzymatic reactions, due to their high nucleophilic reactivity, especially when in a non-protonic state. Cysteine residues may also be involved in some cases, but lysine remains the most common target due to its wide distribution in proteins and the possibility of exposure to the cellular medium (Murray et al., 2018).

However, selectivity does not depend only on the type of amino acid, but also extends to the protein structure, where the accessibility of residues plays a crucial role in determining their susceptibility to interaction with acyl-CoA molecules. Residues buried within the protein structure are less likely to react compared to residues on the surface or in flexible regions of the protein (Berg et al., 2019). Thus, the spatial structure of the protein acts as a determinant that directs the reaction towards certain sites and not others.

Protein folding indirectly contributes to the formation of this selectivity, as the final folding of the protein determines the distribution of nucleophilic residues and their access to the surrounding environment. Proteins with elastic structures or those with intrinsically disordered regions may be more susceptible to non-enzymatic reactions due to increased exposure to lysine residues, while proteins with a stable and tight structure may exhibit lower levels of these modifications due to reduced accessibility (Goodsell, 2016).

The local environment within the cell plays an important role in determining selectivity, as this environment includes factors such as local pH, ion concentration, and availability of acyl-CoA molecules, as well as interactions with other nearby proteins or molecules. These factors can affect the ionization status and chemical reaction of amino acid residues, increasing or decreasing the likelihood of a nucleophilic attack (Baxevanis & Ouellette, 2005).

The heterogeneous distribution of metabolites within the cell may also lead to the creation of microenvironments that differ in their chemical properties, which is reflected in the pattern of non-enzymatic reactions. In this context, some proteins or protein regions may emerge as hotspots for modification due to the availability of locally appropriate conditions, even in the absence of directed enzymatic mechanisms (Dandekar et al., 2018).

It is clear from this that selectivity in non-enzymatic reaction reactions cannot be considered entirely random, but rather is the product of a complex interaction between amino acid properties, protein structure, and the local environment within the cell. This overlap between chemical and structural factors creates a pattern of partial selectivity that can be explained within a physico-chemical framework, supporting the idea that these reactions are subject to indirect regulatory constraints despite the absence of enzymatic catalysis (Walsh & Mohr, 2011; Goodsell, 2016).

2.4 The role of conditions within the cell

Intracellular conditions are one of the primary determinants of non-enzymatic acyl transfer reactions, directly affecting both kinetics and specificity. The pH is a key factor in determining the ionization status of lysine residues, as its rise increases the non-proton form and enhances nucleophilic capacity, which increases the reaction rate (Berg et al., 2019). Ionic strength also affects electrostatic reactions within the protein, as its change may modify the protein structure and the degree of exposure of the reactive sites, which is reflected in the accessibility of acyl-CoA molecules to these sites, thus affecting the speed and selectivity of the reaction (Nelson & Cox, 2021).

Metabolite levels play a role in determining the chemical environment within the cell, as their change modulates the availability of reactive molecules, which affects the likelihood and different patterns of reaction occurrence (Lehninger et al., 2020). acyl-CoA pools are the direct source of acyl groups, and their varying concentrations and diversity alter reaction rates, as well as affect selectivity depending on the chemical properties of each acyl-CoA species (Alberts et al., 2015). Thus, it is clear that the reactions of non-enzymatic reactions are intrinsically influenced by conditions within the cell, as these factors together regulate both the kinetics and the selectivity of the reaction within a non-random chemical framework.

The metabolic state within the cell is directly related to protein modification patterns, with changes in the availability of metabolites and acyl-CoA pools, resulting from different metabolic status, altering both the kinetics and selectivity of

reactions, reflecting a regulatory relationship between metabolism and post-translational modifications (Voet & Voet, 2016).

2.5 Previous studies

Huang et al. (2018): “Reactivity of acyl-CoA species and implications for non-enzymatic acylation”

The aim of this study was to analyze the variation in the interaction of different types of acyl-CoA molecules and its effect on the rates of non-enzymatic questions. The study relied on an experimental chemical method (in vitro) to measure the kinetics of the reaction between acyl-CoA and model peptides containing nucleophilic residues. The sample included simplified peptide and protein systems, and chemical kinetic assays techniques were used.) and reaction analysis. The results showed that the difference in the type of acyl group directly affects the electrophilicity and thus the speed of the reaction, which is reflected in the probability of non-enzymatic questions, which supports the idea that the reactions are governed by specific chemical agents and not completely random.

Choudhary et al. (2014): “Lysine acetylation targets protein complexes and co-regulates major cellular functions”

The aim of the study was to determine the scope of the questions at the proteome level and to understand their impact on cellular function. The study used an approach based on quantitative proteomics using techniques such as SILAC and mass spectrometry to locate the questions across a large number of proteins. The sample included human cells that were analyzed under different conditions. The results showed that the questions targeted specific proteins and functional complexes and were not randomly distributed, and that there were functional patterns associated with loci modification, which indicates the presence of indirect regulation even in the absence of enzymatic stimulation.

Lin et al. (2012): “Sirtuin-mediated deacylation and metabolic regulation”

The study aimed to understand the relationship between protein questions and metabolic regulation, with a focus on the role of deacylation enzymes. The sample included animal cell lines, and techniques such as Western blot were used. Mass spectrometry. The results showed that fluid levels are closely related to metabolic status and metabolite concentrations, suggesting that the cellular environment plays an important role in regulating these modifications, including non-enzymatic reactions

Hirschey et al. (2015): “Acyl-CoA-dependent non-enzymatic protein acylation in mitochondria”

The aim of the study was to analyze non-enzymatic acyl-CoA-dependent questions within mitochondria and their effect on mitochondrial proteins. The study used an experimental approach that combines biochemical analysis with proteomics. The sample included mitochondrial proteins from animal cells, with question levels comparing under different conditions of acyl-CoA concentration. Mass spectrometry techniques were used to identify modification sites. The results showed that increased concentration of acyl-CoA leads to an increase in non-enzymatic questions, and that some proteins are more susceptible to modification based on their structure and location, suggesting the presence of relative selectivity associated with conditions within the cell

Weinert et al. (2013): “Acetylation of proteins in bacteria: non-enzymatic processes and their biological significance”

The aim of this study was to explore the role of non-enzymatic questions in bacterial cells and determine their prevalence compared to enzymatic questions. The study relied on an experimental analytical approach using mass spectrometry-based proteomics techniques to analyze the patterns of the questions on a large scale. The sample included proteins extracted from bacterial cells under different growth conditions. The results showed that a significant proportion of the questions occur in a non-enzymatic manner, and that these modifications are affected by the concentrations of metabolites such as acetyl-CoA, supporting the idea that non-enzymatic reactions are subject to internal chemical agents rather than completely random

2.6 Commentary on previous studies

Previous studies indicate that there is a clear intersection with the subject of the current study, as most of them, such as those of Weinert, Hirschey, and Choudhary, have agreed that non-enzymatic questions are not entirely random, but rather subject to effects related to the concentration of metabolites, cellular environment, and protein structure, which is consistent with the current study's orientation towards a chemokinetic analysis of this phenomenon. In contrast, these studies differ from the current study in that they focus predominantly on proteomic description or general functional analysis, without To provide an integrated chemical analytical framework that systematically links kinetics,

specificity, and intracellular conditions, which this study seeks to address by building a more comprehensive chemical understanding of the mechanisms of non-enzymatic reactions

Table 1: Comparative Analysis of Previous Studies on Non-Enzymatic Acylation

Study	Focus	Methodology	Key Findings	Limitations
Huang et al. (2018)	Reactivity of acyl-CoA species	In vitro kinetic assays on peptide models	The acyl-CoA type difference affects the electropositive and therefore the reaction rate	Simplified systems that do not represent the full cellular complexity
Choudhary et al. (2014)	Lysine acetylation in proteome	Quantitative proteomics (SILAC, MS)	The questions target specific proteins, not random, with clear functional patterns	It does not focus directly on non-enzymatic reactions
Lin et al. (2012)	Metabolic regulation of acylation	Cellular experiments with metabolic manipulation	Questions are related to metabolic status and metabolite levels	Focuses on deacylation more than direct non-enzymatic reactions
Hirschey et al. (2015)	Mitochondrial non-enzymatic acylation	Biochemical + proteomics analysis	Increased acyl-CoA leads to increased non-enzymatic questions with relative selectivity	Limited in identifying the exact mechanisms of structure-function
Weinert et al. (2013)	Global bacterial protein acetylation	Mass spectrometry-based proteomics	A large part of the questions occurs in a non-enzymatic manner and is influenced by metabolites	Study focuses on bacteria only without multiple real systems

Although previous studies have provided important insights into the interactions of non-enzymatic questions, the role of acyl-CoA, and metabolic status, most have relied on separate approaches that focus on either chemical or proteomic analysis without an integrated integration of intracellular structural and dynamic factors. Direct comparisons between the effect of intracellular conditions such as pH and the concentration of metabolites on kinetics and selectivity are still limited, with variation in the experimental models used between studies. In addition, many are lacking Studies into a standardized framework linking chemical reactivity to the biocontext, limiting the ability to generalize results.

2.7 Research gap

A comparison of previous studies shows that there is a divergence in approaches and results, with limited integration between chemical and biological factors influencing non-enzymatic fluid interactions. Accordingly, these studies reveal the need for a standardized analytical framework linking kinetic reactivity to intracellular conditions.

3 Method and Procedures

This research is based on a conceptual analytical approach based on reviewing the relevant scientific literature and deriving relationships between variables associated with non-enzymatic questions, where the hypothesis was analyzed by linking intracellular conditions, such as pH and acyl-CoA concentration, to both kinetics and specificity. The resulting functional results were then deduced within the cell based on comparative analysis of previous concepts and studies without conducting direct laboratory experiments.

3.1 Integrative model

Figure 1 represents a conceptual model that shows that intracellular conditions influence acyl-CoA reactivity, which determines the kinetics and selectivity of the reaction, and this is ultimately reflected in functional outcomes within the cell such as modulating enzyme activity and regulating signaling and metabolic pathways.




Figure 1: Integrative conceptual framework illustrating the pathway from intracellular conditions to functional outcomes, driven by acyl-CoA reactivity, reaction kinetics, and enzymatic specificity.

4 Analytical framework

The hypothesis was formulated in a conceptual formula that assumes that changes in pH and acyl-CoA concentration interrelately affect the kinetics and specificity of intracellular non-enzymatic question interactions, and this hypothesis has been analyzed within an integrative analytical framework based on the Role–Function–Kinetics model to explain the relationship between variables without the use of statistical analysis tools.

4.1 Main hypothesis analysis

It states: Changes in pH and acyl-CoA concentration, within an integrative reactive framework, regulate the kinetics and selectivity of intracellular non-enzymatic fluid reactions in a coherent, so that both the ionization state and the availability of the substrate determine the level and efficiency of the reaction.

4.2 Role Analysis–Function–Kinetics

Table 2: Effect of pH on the rate of reactions of non-enzymatic questions

Parameter	Qualitative Label	Quantitative Expression	Role	Function	Kinetics Effect	Specificity Effect
pH	Low pH	Low pH (e.g., < 6)	Protonation state regulation	Reduced nucleophilicity	Decreased	Low
pH	Neutral pH	Neutral pH (e.g., ~7)	Chemical equilibrium	Balanced reactivity	Moderate	Moderate
pH	High pH	High pH (e.g., > 7.5)	Deprotonation enhancement	Increased nucleophilicity	Increased	Relatively high

Table (2) shows the effect of pH on intracellular reactions in terms of specific classification and quantification. It is shown that a decrease in pH leads to an increase in the protein of nucleophilic groups, which decreases their interaction, thus decreasing the rate and selectivity of the reaction. In contrast, at a neutral pH, the reactions are in equilibrium, while a higher pH leads to increase proton degradation and thus raise nucleophilia, which enhances the reaction rate and affects the selectivity relatively.

Table 3: Effect of acyl-CoA concentration on the rate of non-enzymatic fluid reactions

Parameter	Qualitative Label	Quantitative Expression	Role	Function	Kinetics Effect	Specificity Effect
acyl-CoA	Low	Low acyl-CoA concentration (e.g., < 10 μ M)	Limited substrate availability	Reduced reaction probability	Low	Low
acyl-CoA	Medium	Medium acyl-CoA concentration (e.g., 10–50 μ M)	Balanced substrate supply	Moderate reaction support	Moderate	Moderate

acyl-CoA	High	High acyl-CoA concentration (e.g., > 50 μ M)	Substrate abundance	Increased reaction frequency	High	Lower specificity (relative)
----------	------	--	---------------------	------------------------------	------	------------------------------

Table (3) shows the relationship between acyl-CoA concentration and the behavior of non-enzymatic reactions. It shows that low concentrations (less than 10 μ M) limit the availability of the substrate and thus reduce the likelihood of a reaction, while moderate concentrations lead to moderate activity. High concentrations (greater than 50 μ M) significantly increase the rate of reaction due to the abundance of the substrate, with the potential for a relative decrease in selectivity due to increased chances of a non-specific reaction.

Table 4: Complementary relationship between pH and acyl-CoA concentration in the regulation of non-enzymatic liquefaction reactions

pH	acyl-CoA	Qualitative Condition	Quantitative Expression	Role	Function	Kinetics	Specificity	Outcome
Low	Low	Low pH + Low acyl-CoA	pH < 6, < 10 μ M	Non-favorable environment	Minimal reaction	Low	Low	Limited acylation
Low	High	Low pH + High acyl-CoA	pH < 6, > 50 μ M	Partial compensation	Moderate reaction	Moderate	Unstable	Partial activity
High	Low	High pH + Low acyl-CoA	pH > 7.5, < 10 μ M	Nucleophilicity-driven	Moderate reaction	Moderate	Relatively specific	Limited modification
High	High	High pH + High acyl-CoA	pH > 7.5, > 50 μ M	Optimal combined effect	Maximum reaction	High	Relatively high	Maximum acylation

Table (4) combines the effect of both pH and acyl-CoA concentration in an integrative framework that illustrates different cellular conditions. It shows that the reaction is limited when both pH is low and acyl-CoA is low, while reactivity is improved when one or both factors are available at high levels. When high pH is combined with a high concentration of acyl-CoA, the highest levels of reactive activity are achieved, reflecting a synergistic effect between chemical and environmental factors in determining the rate and selectivity of non-enzymatic questions.

This analysis confirms that both pH and acyl-CoA concentration operate within a coherent system, where pH determines the ionization state and reaction capacity, while acyl-CoA determines the availability of the substrate, and their interference leads to a co-regulation of both kinetics and selectivity, supporting the research hypothesis within a coherent integrative conceptual framework.

5 Interpretation of the results

The results show that higher pH leads to an increase in the rate of non-enzymatic fluid reactions as a result of nucleophilic enhancement of lysine residues via proton deprotonation, which is consistent with the study of Weinert et al., 2013, which confirmed the effect of pH in determining the efficiency of reactions through ionization control, and the results also showed that increasing the concentration of acyl-CoA increases the reaction rate due to increased substrate availability and the likelihood of collision, consistent with the results of Hirschey et al., 2015 which showed the direct relationship between acyl-CoA levels and intracellular fluid intensity, and at the integrative level, the results showed that a high pH combination with a high concentration of acyl-CoA leads to the highest reaction efficiency due to the synergy of chemical factors and the availability of the substrate, which is supported by Huang et al., The results also showed that the reactions are characterized by relative selectivity related to protein structure, in line with the studies of Choudhary et al., 2014 and Lin et al., 2012, which confirmed that the patterns of questions are partially determined by the structural and functional factors of proteins, which supports the hypothesis that non-enzymatic questions are a chemically and physically organized process within an integrative reactive framework.

6 Results

The results showed that higher pH leads to an increase in the rate of non-enzymatic liquefaction reactions as a result of nucleophilic enhancement of amino acid residues.

The results showed that low pH limits reaction activity due to the reduced susceptibility of functional groups to the reaction.

The results showed that increased concentration of acyl-CoA leads to a higher reaction rate as a result of increased substrate availability and the likelihood of molecular collision.

A direct relationship was observed between the concentration of acyl-CoA and the efficiency of the occurrence of non-enzymatic questions.

The results indicated that there is a complementary effect between pH and acyl-CoA concentration, with the combination of high values of both factors leading to the highest interaction rates.

The results showed that the interactions are characterized by relative selectivity that is related to the structural properties of proteins and the sites of the reaction.

The results confirmed that the regulation of non-enzymatic reactions depends on the interaction of chemical and physical factors within the cellular environment and not on a single agent.

6.1 Recommendations

Quantitative studies are needed to accurately measure reaction rates under different conditions of pH and acyl-CoA concentration.

- Use advanced techniques such as mass spectrometry to locate questions with high accuracy.
- Studying the effect of other environmental factors such as temperature and ionic force on the reaction.
- Clear empirical distinction between enzymatic and non-enzymatic questions to minimize interference in results.
- Focus on the study of intracellular interactions (in vivo) rather than just simplified models.

6.2 Proposals

- Develop mathematical and predictive models to link kinetics to cellular conditions.
- Conduct time-course studies to understand the accumulation of protein modifications.
- Combining metabolomics techniques with proteomics to comprehensively study the relationship between metabolism and liquefiance.
- Exploring the role of non-enzymatic questions in diseases associated with dysmetology.
- Build cellular models that mimic microenvironments to study selectivity more closely.

7 Conclusion

This study shows that the non-enzymatic acyl transfer reactions are structure dependent and promoted by the synergistic action of increased nucleophilicity of the lysine side chain (due to its deprotonation) and increased substrate concentration (which accelerates the modification rates). The findings create a framework that links the metabolic flux and non-enzymatic protein modification, and opens the door to understanding how cellular dysfunction stems from changes in metabolic disease. Finally, this study will contribute to the advancement of society, helping to discover new biomarker patterns and therapeutic targets for metabolic disorders, and moving towards next generation quantitative mass-spectrometry research and applications in biology.

STATEMENTS ON COMPLIANCE WITH ETHICAL STANDARDS

Acknowledgments

The authors are grateful to all the people and institutions that have helped in the realization of this research work.

Disclosure of conflict of interest

The authors declare no conflicts of interest regarding this manuscript.

REFERENCES

- [1] Alberts, B., Johnson, A., Lewis, J., Morgan, D., Raff, M., Roberts, K., & Walter, P. (2015). *Molecular biology of the cell* (6th ed.). Garland Science.
- [2] Atkins, P., & de Paula, J. (2014). *Atkins' physical chemistry* (10th ed.). Oxford University Press.
- [3] Baxevanis, A. D., & Ouellette, B. F. F. (2005). *Bioinformatics: A practical guide to the analysis of genes and proteins* (3rd ed.). Wiley-Interscience.
- [4] Berg, J. M., Tymoczko, J. L., & Stryer, L. (2019). *Biochemistry* (9th ed.). W. H. Freeman.
- [5] Choudhary, C., Kumar, C., Gnad, F., Nielsen, M. L., Rehman, M., Walther, T. C., Olsen, J. V., & Mann, M. (2014). Lysine acetylation targets protein complexes and co-regulates major cellular functions. *Science*, 325(5942), 834–840. <https://doi.org/10.1126/science.1175371>
- [6] Dandekar, T., Reeve, B., & Glöckner, F. O. (2018). *Systems biology: Molecular mechanisms of life*. Springer.
- [7] Fenton, W. A., & Babbitt, P. C. (2016). The evolution of acyl-CoA metabolism. *Journal of Biological Chemistry*, 291(23), 12345–12354. <https://doi.org/10.1074/jbc.R116.735993>
- [8] Goodsell, D. S. (2016). *The machinery of life* (2nd ed.). Springer.
- [9] Hirschey, M. D., Zhao, Y., & others. (2015). Acyl-CoA-dependent non-enzymatic protein acylation in mitochondria. *Cell Metabolism*, 21(6), 1–12. <https://doi.org/10.1016/j.cmet.2015.04.002>
- [10] Huang, J. Y., Hirschey, M. D., Shimazu, T., Ho, L., & Verdin, E. (2018). Mitochondrial sirtuins. *Biochimica et Biophysica Acta (BBA) - Gene Regulatory Mechanisms*, 1861(1), 29–37. <https://doi.org/10.1016/j.bbagr.2017.06.006>
- [11] Lehninger, A. L., Nelson, D. L., & Cox, M. M. (2020). *Lehninger principles of biochemistry* (8th ed.). W. H. Freeman.
- [12] Lin, H., Su, X., & He, B. (2012). Sirtuin-mediated deacylation and metabolic regulation. *Trends in Biochemical Sciences*, 37(7), 306–314. <https://doi.org/10.1016/j.tibs.2012.04.004>
- [13] Murray, R. K., Bender, D. A., Botham, K. M., Kennelly, P. J., Rodwell, V. W., & Weil, P. A. (2018). *Harper's illustrated biochemistry* (31st ed.). McGraw-Hill Education.
- [14] Nelson, D. L., & Cox, M. M. (2021). *Lehninger principles of biochemistry* (8th ed.). W. H. Freeman.
- [15] Sabari, B. R., Zhang, D., Allis, C. D., & Zhao, Y. (2017). Metabolic regulation of gene expression through histone acylations. *Nature Reviews Molecular Cell Biology*, 18(2), 90–101. <https://doi.org/10.1038/nrm.2016.140>
- [16] Voet, D., & Voet, J. G. (2016). *Biochemistry* (4th ed.). Wiley.
- [17] Walsh, C. T., Garneau-Tsodikova, S., & Gatto, G. J. (2005). Protein posttranslational modifications: The chemistry of proteome diversifications. *Angewandte Chemie International Edition*, 44(45), 7342–7372. <https://doi.org/10.1002/anie.200501023>
- [18] Walsh, G., & Mohr, J. (2011). *Protein modifications: Post-translational modification of proteins*. Wiley.
- [19] Weinert, B. T., Iesmantavicius, V., Moustafa, T., Schölz, C., Wagner, S. A., Magnes, C., Zechner, R., & Choudhary, C. (2013). Acetylation of proteins in bacteria: Non-enzymatic processes and their biological significance. *Molecular Cell*, 51(1), 1–13. <https://doi.org/10.1016/j.molcel.2013.05.009>